

**Clinical Evidence Supporting the Need for Glycemic Control in the Hospital Setting
“The Case for Glycemic Control in Hospitalized Patients”**

References and Key Readings

Most Recent Consensus Statements

Moghissi ES, Korytkowski MT, DiNardo M; American Association of Clinical Endocrinologists; American Diabetes Association. American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycemic control. *Endocr Pract.* 2009;15:353-69.

Moghissi ES, Korytkowski MT, DiNardo M; American Association of Clinical Endocrinologists; American Diabetes Association. American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycemic control. *Diabetes Care.* 2009;32:1119-31.

2009

American Diabetes Association. Standards of Medical Care in Diabetes—2009. *Diabetes Care* 2009;32:S13-S61.

Bailon RM, Partlow BJ, Miller-Cage V, et al. Continuous subcutaneous insulin infusion (insulin pump) therapy can be safely used in the hospital in select patients. *Endocr Pract.* 2009;15:24-9.

Hammer MJ, Casper C, Gooley TA, et al. The contribution of malglycemia to mortality among allogeneic hematopoietic cell transplant recipients. *Biol Blood Marrow Transplant.* 2009;15:344-51.

Hirsch IB. Sliding scale insulin--time to stop sliding. *JAMA.* 2009;301:213-4.

Inzucchi SE, Siegel MD. Glucose control in the ICU – how tight is too tight? *N Engl J Med* 2009;360:1346-9.

Korytkowski MT, Salata RJ, Koerbel GL, et al. Insulin Therapy and Glycemic Control in Hospitalized Patients With Diabetes During Enteral Nutrition Therapy. *Diabetes Care* 2009; 32:594–596.

Kosiborod M, Inzucchi SE, Goyal A, et al. Relationship between spontaneous and iatrogenic hypoglycemia and mortality in patients hospitalized with acute myocardial infarction. *JAMA.* 2009;301:1556-64.

Mowery NT, Gunter OL, Guillaumondegui O, et al. Stress insulin resistance is a marker for mortality in traumatic brain injury. *J Trauma.* 2009;66:145-51; discussion 151-3.

NICE-SUGAR Study Investigators. Intensive versus conventional glucose control in critically ill patients. *N Engl J Med* 2009;360:1283-97.

Scott MG, Bruns DE, Boyd JC, et al. Tight glucose control in the intensive care unit: are glucose meters up to the task? *Clin Chem*. 2009;55:18-20.

Seley J, Wallace M. Meeting the Challenge of Inpatient Diabetes Education: An Interdisciplinary Approach. *Educating Your Patient with Diabetes*. P. 81-96.

Umpierrez GE. Basal versus sliding-scale regular insulin in hospitalized patients with hyperglycemia during enteral nutrition therapy. *Diabetes Care* 2009; 32:751-3.

Umpierrez GE, Hor T, Smiley D, et al. Comparison of Inpatient Insulin Regimens with Detemir plus Aspart Versus Neutral Protamine Hagedorn plus Regular in Medical Patients with Type 2 Diabetes. *J Clin Endocrinol Metab*. 2009;94:564-9.

van der Crabben SN, Blümer RM, Stegenga ME, et al. Early endotoxemia increases peripheral and hepatic insulin sensitivity in healthy humans. *J Clin Endocrinol Metab*. 2009;94:463-8.

Vlasselaers D, Milants I, Desmet L, et al. Intensive insulin therapy for patients in paediatric intensive care: a prospective, randomised controlled study. *Lancet*. 2009;373:547-56.

2008

American Diabetes Association: Economic costs of diabetes in the U.S. in 2007. *Diabetes Care* 2008;31:596-615.

Amori RE, Pittas AG, Siegel RD, et al. Inpatient medical errors involving glucose-lowering medications and their impact on patients: review of 2,598 incidents from a voluntary electronic error-reporting database. *Endocr Pract*. 2008;14:535-42.

Arinzon Z, Shabat S, Shuval I, et al. Prevalence of diabetes mellitus in elderly patients received enteral nutrition long-term care service. *Arch Gerontol Geriatr*. 2008;47:383-93.

Beardsall K, Vanhaesebrouck S, Ogilvy-Stuart AL, et al. Early insulin therapy in very-low-birth-weight infants. *N Engl J Med*. 2008;359:1873-84.

Brunkhorst FM, Engel C, Bloos F, for the German Competence Network Sepsis (SepNet). Intensive insulin therapy and pentastarch resuscitation in severe sepsis. *N Engl J Med*. 2008;358:125-39.

Bruno A, Gregorio D, Caropreso A. Normal glucose values are associated with a lower risk of mortality in hospitalized patients. *Diabetes Care* 2008;31:2209-2210.

Centers for Disease Control and Prevention. National diabetes fact sheet: general information and national estimates on diabetes in the United States, 2007. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, 2008.

Cochran A, Davis L, Morris SE, et al. Safety and efficacy of an intensive insulin protocol in a burn-trauma intensive care unit. *J Burn Care Res*. 2008;29:187-91.

Cook CB, Jameson KA, Hartsell ZC, et al. Beliefs about hospital diabetes and perceived barriers to glucose management among inpatient midlevel practitioners. *Diabetes Educ.* 2008;34:75-83.

De La Rosa GC, Donado JH, Restrepo AH: GICIHPTU: Strict glycaemic control in patients hospitalised in a mixed medical and surgical intensive care unit: a randomised clinical trial. *Critical Care* 2008,12:R120.

Deedwania P, Kosiborod M, Barrett E; American Heart Association Diabetes Committee of the Council on Nutrition, Physical Activity, and Metabolism. Hyperglycemia and acute coronary syndrome: a scientific statement from the American Heart Association Diabetes Committee of the Council on Nutrition, Physical Activity, and Metabolism. *Circulation* 2008;117:1610-9.

Derr RL, Hsiao VC, Saudek CD. Antecedent hyperglycemia is associated with an increased risk of neutropenic infections during bone marrow transplantation. *Diabetes Care* 2008;31:1972-7.

Duane TM, Ivatury RR, Dechert T, et al. Blood glucose levels at 24 hours after trauma fails to predict outcomes. *J Trauma.* 2008;64:1184-7.

Falcao G, Ulate K, Kouzekanani K, et al. Impact of postoperative hyperglycemia following surgical repair of congenital cardiac defects. *Pediatr Cardiol.* 2008;29:628-36.

Finfer S, Delaney A. Tight glycaemic control in critically ill adults. *JAMA.* 2008;300:963-5.

Kitabchi AE, Freire AX, Umpierrez GE. Evidence for strict inpatient blood glucose control: time to revise glycaemic goals in hospitalized patients. *Metabolism.* 2008;57:116-20.

Kosiborod M, Inzucchi SE, Krumholz HM, et al. Glucometrics in patients hospitalized with acute myocardial infarction: defining the optimal outcomes-based measure of risk. *Circulation.* 2008;117:1018-27.

Matheny ME, Shubina M, Kimmel ZM, et al. Treatment intensification and blood glucose control among hospitalized diabetic patients. *J Gen Intern Med.* 2008;23:184-9.

Mills RD, Schwartz F, Shubrook JH. Evaluation of diabetes management in a rural community hospital. *Endocr Pract.* 2008;14:50-5.

Moghissi ES. Insulin strategies for managing inpatient and outpatient hyperglycemia and diabetes. *Mt Sinai J Med.* 2008;75:558-66.

Noschese M, Donihi AC, Koerbel G, et al. Effect of a diabetes order set on glycaemic management and control in the hospital. *Qual Saf Health Care.* 2008;17:464-8.

Palacio A, Smiley D, Ceron M, et al. Prevalence and clinical outcome of inpatient hyperglycemia in a community pediatric hospital. *J Hosp Med.* 2008;3:212-7.

Pasternak JJ, McGregor DG, Schroeder DR; IHAST Investigators. Hyperglycemia in patients undergoing cerebral aneurysm surgery: its association with long-term gross

neurologic and neuropsychological function. *Mayo Clin Proc.* 2008;83:406-17.

Saudek CD, Herman WH, Sacks DB, et al. A new look at screening and diagnosing diabetes mellitus. *J Clin Endocrinol Metab.* 2008;93:2447-53.

Treggiari MM, Karir V, Yanez ND, et al. Intensive insulin therapy and mortality in critically ill patients. *Crit Care.* 2008;12:R29.

Trujillo JM, Barsky EE, Greenwood BC, et al. Improving glycemic control in medical inpatients: a pilot study. *J Hosp Med.* 2008;3(1):55-63.

Tuggle DW, Kuhn MA, Jones SK, et al. Hyperglycemia and infections in pediatric trauma patients. *Am Surg.* 2008;74:195-8.

Wahl WL, Taddonio M, Maggio PM, et al. Mean glucose values predict trauma patient mortality. *J Trauma.* 2008;65:42-7.

Wiener RS, Wiener DC, Larson RJ. Benefits and risks of tight glucose control in critically ill adults: a meta-analysis. *JAMA* 2008;300:933-944.

2007

Anthony M. Treatment of hypoglycemia in hospitalized adults: a descriptive study. *Diabetes Educ.* 2007;33:709-15.

Asudani D, Calles-Escandon J. Inpatient hyperglycemia: slide through the scale but cover the bases first. *J Hosp Med.* 2007;2 Suppl 1:23-32.

Becker T, Moldoveanu A, Cukierman T, et al. Clinical outcomes associated with the use of subcutaneous insulin-by-glucose sliding scales to manage hyperglycemia in hospitalized patients with pneumonia. *Diabetes Res Clin Pract.* 2007;78:392-7.

Bochicchio GV, Joshi M, Bochicchio KM, et al. Early hyperglycemic control is important in critically injured trauma patients. *J Trauma.* 2007;63:1353-8.

Clement S. Better glycemic control in the hospital: beneficial and feasible. *Cleve Clin J Med.* 2007;74:111-2, 114-20.

Cohen LS, Sedhom L, Salifu M, et al. Inpatient diabetes management: examining morning practice in an acute care setting. *Diabetes Educ.* 2007;33:483-92.

Cook CB, Castro JC, Schmidt RE, et al. Diabetes care in hospitalized noncritically ill patients: More evidence for clinical inertia and negative therapeutic momentum. *J Hosp Med.* 2007;2:203-11.

Cook CB, Moghissi E, Joshi R, et al. Inpatient point-of-care bedside glucose testing: preliminary data on use of connectivity informatics to measure hospital glycemic control. *Diabetes Technol Ther.* 2007;9:493-500.

Cook CB, McNaughton DA, Braddy CM, et al. Management of inpatient hyperglycemia:

assessing perceptions and barriers to care among resident physicians. *Endocr Pract.* 2007;13:117-24.

Cook CB, Stockton L, Baird M, et al. Working to improve care of hospital hyperglycemia through statewide collaboration: the Georgia Hospital Association Diabetes Special Interest Group. *Endocr Pract.* 2007;13:45-50.

Dungan K, Chapman J, Braithwaite SS, et al. Glucose measurement: confounding issues in setting targets for inpatient management. *Diabetes Care.* 2007;30:403-9.

Elinav H, Wolf Z, Szalat A, et al. In-hospital treatment of hyperglycemia: effects of intensified subcutaneous insulin treatment. *Curr Med Res Opin.* 2007;23:757-65.

Fuji S, Kim SW, Mori S, et al. Hyperglycemia during the neutropenic period is associated with a poor outcome in patients undergoing myeloablative allogeneic hematopoietic stem cell transplantation. *Transplantation.* 2007;84:814-20.

Gale SC, Sicoutris C, Reilly PM, et al. Poor glycemic control is associated with increased mortality in critically ill trauma patients. *Am Surg.* 2007;73:454-60.

Gandhi GY, Nuttall GA, Abel MD, et al. Intensive intraoperative insulin therapy versus conventional glucose management during cardiac surgery: a randomized trial. *Ann Intern Med.* 2007;146:233-43.

Garg R, Bhutani H, Jarry A, et al. Provider response to insulin-induced hypoglycemia in hospitalized patients. *J Hosp Med.* 2007;2:258-60.

Hassan E. Hyperglycemia management in the hospital setting. *Am J Health Syst Pharm.* 2007;64(10 Suppl 6):S9-S14.

Held C, Gerstein HC, Yusuf S; ONTARGET/TRANSCEND Investigators. Glucose levels predict hospitalization for congestive heart failure in patients at high cardiovascular risk. *Circulation.* 2007;115:1371-5.

Hovorka R, Kremen J, Blaha J, et al. Blood glucose control by a model predictive control algorithm with variable sampling rate versus a routine glucose management protocol in cardiac surgery patients: a randomized controlled trial. *J Clin Endocrinol Metab.* 2007;92:2960-4.

Karon BS, Gandhi GY, Nuttall GA, et al. Accuracy of roche accu-chek inform whole blood capillary, arterial, and venous glucose values in patients receiving intensive intravenous insulin therapy after cardiac surgery. *Am J Clin Pathol.* 2007;127:919-26.

Kim S. Burden of hospitalizations primarily due to uncontrolled diabetes: implications of inadequate primary health care in the United States. *Diabetes Care.* 2007;30:1281-2.

Krinsley JS, Grover A. Severe hypoglycemia in critically ill patients: risk factors and outcomes. *Crit Care Med.* 2007;35:2262-7.

Kripalani S, Jackson AT, Schnipper JL, et al. Promoting effective transitions of care at

hospital discharge: a review of key issues for hospitalists. *J Hosp Med.* 2007;2:314-23.

Lansang MC, Harrell H. Knowledge on inpatient diabetes among fourth-year medical students. *Diabetes Care.* 2007;30:1088-91.

Meynaar IA, Dawson L, Tangkau PL, et al. Introduction and evaluation of a computerised insulin protocol. *Intensive Care Med.* 2007;33:591-6.

Nazer LH, Chow SL, Moghissi ES. Insulin infusion protocols for critically ill patients: a highlight of differences and similarities. *Endocr Pract.* 2007;13:137-46.

Noordzij PG, Boersma E, Schreiner F, et al. Increased preoperative glucose levels are associated with perioperative mortality in patients undergoing noncardiac, nonvascular surgery. *Eur J Endocrinol* 2007;156:137-142.

Rea RS, Donihi AC, Bobeck M, et al. Implementing an intravenous insulin infusion protocol in the intensive care unit. *Am J Health Syst Pharm.* 2007;64:385-95.

Reed CC, Stewart RM, Sherman M, et al. Intensive insulin protocol improves glucose control and is associated with a reduction in intensive care unit mortality. *J Am Coll Surg.* 2007;204:1048-54; discussion 1054-5.

Reynolds LR, Cook AM, Lewis DA, et al. An institutional process to improve inpatient glycemic control. *Qual Manag Health Care.* 2007;16:239-49.

Rodbard HW, Blonde L, Braithwaite SS; AACE Diabetes Mellitus Clinical Practice Guidelines Task Force. American Association of Clinical Endocrinologists medical guidelines for clinical practice for the management of diabetes mellitus. *Endocr Pract.* 2007 May-Jun;13 Suppl 1:1-68.

Rubin DJ, Moshang J, Jabbour SA. Diabetes knowledge: are resident physicians and nurses adequately prepared to manage diabetes? *Endocr Pract.* 2007;13:17-21.

Shin S, Britt RC, Reed SF, et al. Early glucose normalization does not improve outcome in the critically ill trauma population. *Am Surg.* 2007;73:769-72.

Umpierrez GE. Inpatient management of diabetes: an increasing challenge to the hospitalist physician. *J Hosp Med.* 2007;2 Suppl 1:33-5.

Umpierrez GE, Smiley D, Zisman A, et al. Randomized study of basal-bolus insulin therapy in the inpatient management of patients with type 2 diabetes (RABBIT 2 trial). *Diabetes Care.* 2007;30:2181-6.

Umpierrez GE, Palacio A, Smiley D. Sliding scale insulin use: myth or insanity? *Am J Med.* 2007;120:563-7.

Wexler DJ, Meigs JB, Cagliero E, et al. Prevalence of hyper- and hypoglycemia among inpatients with diabetes: a national survey of 44 U.S. hospitals. *Diabetes Care.* 2007;30:367-9.

Wheeler K, Crawford R, McAdams D, et al. Inpatient to outpatient transfer of diabetes care: perceptions of barriers to postdischarge followup in urban African American patients. *Ethn Dis*. 2007;17:238-43.

2006

American College of Endocrinology Task Force on Inpatient Diabetes and Glycemic. ACE Position Statement on Inpatient Diabetes and Metabolic Control. *Endocr Pract*. 2006;12:3-13.

Baker EH, Janaway CH, Philips BJ, et al. Hyperglycaemia is associated with poor outcomes in patients admitted to hospital with acute exacerbations of chronic obstructive pulmonary disease. *Thorax*. 2006;61:284-289.

Cheung NW, Wong VW, McLean M. The Hyperglycemia: Intensive Insulin Infusion in Infarction (HI-5) study: a randomized controlled trial of insulin infusion therapy for myocardial infarction. *Diabetes Care*. 2006;29:765-770.

Cryer PE. Hypoglycaemia: the limiting factor in the glycaemic management of the critically ill? *Diabetologia*. 2006;49:1722-5.

DeSantis AJ, Schmeltz LR, Schmidt K, et al. Inpatient management of hyperglycemia: the Northwestern experience. *Endocr Pract*. 2006;12:491-505.

Donihi AC, Raval D, Saul M, et al. Prevalence and predictors of corticosteroid-related hyperglycemia in hospitalized patients. *Endocr Pract*. 2006;12:358-362.

Egi M, Bellomo R, Stachowski E, et al. Variability of blood glucose concentration and short-term mortality in critically ill patients. *Anesthesiology*. 2006;105:244-52.

Fonseca V. Newly diagnosed diabetes/hyperglycemia in hospitals: what should we do? *Endocr Pract*. 2006;12 Suppl 3:108-11.

Frontera JA, Fernandez A, Claassen J, et al. Hyperglycemia after SAH: predictors, associated complications, and impact on outcome. *Stroke*. 2006;37:199-203.

Furnary AP, Wu Y. Clinical effects of hyperglycemia in the cardiac surgery population: the Portland Diabetic Project. *Endocr Pract*. 2006;12 Suppl 3:22-6.

Goldberg PA, Bozzo JE, Mesmer M, et al. "Glucometrics" - Assessing the quality of inpatient glucose management. *Diab Technol Therap* 2006;8:560-9.

Goyal A, Mahaffey KW, Garg J, et al. Prognostic significance of the change in glucose level in the first 24 h after acute myocardial infarction: results from the CARDINAL study. *Eur Heart J*. 2006;27:1289-97.

Green Conaway DL, Enriquez JR, Barberena JE, et al. Assessment of and physician response to glycemic control in diabetic patients presenting with an acute coronary syndrome. *Am Heart J*. 2006;152:1022-7.

Hellman R. Patient safety and inpatient glycemic control: translating concepts into action. *Endocr Pract*. 2006 Jul-Aug;12 Suppl 3:49-55.

Hirsch IB. Inpatient diabetes: review of data from the cardiac care unit. *Endocr Pract.* 2006;12 Suppl 3:27-34.

Knecht LA, Gauthier SM, Castro JC, et al. Diabetes care in the hospital: is there clinical inertia? *J Hosp Med.* 2006;1:151-60.

Korytkowski M, DiNardo M, Donihi AC, et al. Evolution of a diabetes inpatient safety committee. *Endocr Pract.* 2006;Suppl 3:91-99.

Krinsley JS, Jones RL. Cost analysis of intensive glycemic control in critically ill adult patients. *Chest.* 2006;129:644-50.

Krinsley JS. Glycemic control, diabetic status, and mortality in a heterogeneous population of critically ill patients before and during the era of intensive glycemic management: six and one-half years experience at a university-affiliated community hospital. *Semin Thorac Cardiovasc Surg.* 2006;18:317-25.

Newton CA, Young S. Financial implications of glycemic control: results of an inpatient diabetes management program. *Endocr Pract.* 2006 Jul-Aug;12 Suppl 3:43-8.

Olson L, Muchmore J, Lawrence CB. The benefits of inpatient diabetes care: improving quality of care and the bottom line. *Endocr Pract.* 2006;12 Suppl 3:35-42.

Pittas AG, Siegel RD, Lau J. Insulin therapy and in-hospital mortality in critically ill patients: systematic review and meta-analysis of randomized controlled trials. *JPEN J Parenter Enteral Nutr.* 2006;30:164-72.

Schmeltz LR, Desantis AJ, Schmidt K, et al. Conversion of intravenous insulin infusions to subcutaneously administered insulin glargine in patients with hyperglycemia. *Endocr Pract.* 2006;12:641-650.

Schnipper JL, Barsky EE, Shaykevich S, et al. Inpatient management of diabetes and hyperglycemia among general medicine patients at a large teaching hospital. *J Hosp Med.* 2006;1:145-50.

Shim YH, Kweon TD, Lee JH, et al. Intravenous glucose-insulin- potassium during off-pump coronary artery bypass surgery does not reduce myocardial injury. *Acta Anaesthesiol Scand.* 2006;50:954-61.

Taylor BE, Schallom ME, Sona CS, et al. Efficacy and safety of an insulin infusion protocol in a surgical ICU. *J Am Coll Surg.* 2006;202:1-9.

Umpierrez G, Maynard G. Glycemic chaos (not glycemic control) still the rule for inpatient care: how do we stop the insanity? *J Hosp Med.* 2006;1:141-4.

Van den Berghe G, Wouters PJ, Kesteloot K, et al. Analysis of healthcare resource utilization with intensive insulin therapy in critically ill patients. *Crit Care Med.* 2006;34:612-616.

Van den Berghe G, Wilmer A, Hermans G, et al. Intensive insulin therapy in the medical ICU. *N*

Engl J Med. 2006;354:449-461.

Vanhorebeek I, Langouche L, Van den Berghe G. Intensive insulin therapy in the intensive care unit: update on clinical impact and mechanisms of action. *Endocr Pract.* 2006;12 Suppl 3:14-22.

Wintergerst KA, Buckingham B, Gandrud L, et al. Association of hypoglycemia, hyperglycemia, and glucose variability with morbidity and death in the pediatric intensive care unit. *Pediatrics.* 2006;118:173-9.

2005

Ainla T, Baburin A, Teesalu R, et al. The association between hyperglycaemia on admission and 180-day mortality in acute myocardial infarction patients with and without diabetes. *Diabet Med.* 2005;22:1321-1325.

Baldwin D, Villanueva G, McNutt R, et al. Eliminating inpatient sliding-scale insulin: a reeducation project with medical house staff. *Diabetes Care.* 2005;28:1008-11.

Bates D, Clark NG, Cook RI, et al. American College of Endocrinology and American Association of Clinical Endocrinologists position statement on patient safety and medical system errors in diabetes and endocrinology. *Endocr Pract.* 2005;11:197-202.

Bochicchio GV, Sung J, Joshi M, et al. Persistent hyperglycemia is predictive of outcome in critically ill trauma patients. *J Trauma.* 2005;58:921-924.

Branco RG, Garcia PC, Piva JP, et al. Glucose level and risk of mortality in pediatric septic shock. *Pediatr Crit Care Med.* 2005;6:470-2.

Cheung NW, Napier B, Zaccaria C, et al. Hyperglycemia is associated with adverse outcomes in patients receiving total parenteral nutrition. *Diabetes Care.* 2005;28:2367-2371.

Collier B, Diaz J Jr, Forbes R, et al. The impact of a normoglycemic management protocol on clinical outcomes in the trauma intensive care unit. *JPEN J Parenter Enteral Nutr.* 2005;29:353-8; discussion 359.

Conaway DG, O'Keefe JH, Reid KJ, et al. Frequency of undiagnosed diabetes mellitus in patients with acute coronary syndrome. *Am J Cardiol.* 2005;96:363-365.

Cook CB. Use of Continuous Subcutaneous Insulin Infusion (Insulin Pump) Therapy in the Hospital Setting: Proposed Guidelines and Outcomes Measures. *Diabetes Educ.* 2005;31:849-857.

CREATE-ECLA Trial Group Investigators. Effect of glucose-insulin-potassium infusion on mortality in patients with acute ST-segment elevation myocardial infarction. *JAMA* 2005;293:437-46.

D'Orazio P, Burnett RW, Fogh-Andersen N; International Federation of Clinical Chemistry Scientific Division Working Group on Selective Electrodes and Point of Care Testing. Approved IFCC recommendation on reporting results for blood glucose (abbreviated). *Clin Chem.* 2005;51:1573-6.

Faustino EV, Apkon M. Persistent hyperglycemia in critically ill children. *J Pediatr.* 2005;146:30-4.

Hirsch IB, Brownlee M. Should minimal blood glucose variability become the gold standard of glycemic control? *J Diabetes Complications.* 2005;19:178-81.

Ishihara M, Kojima S, Sakamoto T; Japanese Acute Coronary Syndrome Study Investigators. Acute hyperglycemia is associated with adverse outcome after acute myocardial infarction in the coronary intervention era. *Am Heart J.* 2005;150:814-820.

Jeremitsky E, Omert LA, Dunham CM, et al. The impact of hyperglycemia on patients with severe brain injury. *J Trauma.* 2005;58:47-50.

Kosiborod M, Rathore SS, Inzucchi S et al. Admission glucose and mortality in elderly patients hospitalized with acute myocardial infarction: implications for patients with and without recognized diabetes. *Circulation.* 2005;111:3078-3086.

Lukins MB, Manninen PH. Hyperglycemia in patients administered dexamethasone for craniotomy. *Anesthesia & Analgesia.* 2005;100:1129-1133.

Malmberg K, Ryden L, Wedel H; DIGAMI 2 Investigators. Intense metabolic control by means of insulin in patients with diabetes mellitus and acute myocardial infarction (DIGAMI 2): effects on mortality and morbidity. *Eur Heart J.* 2005;26:650-661.

McAlister FA, Majumdar SR, Blitz S, et al. The relation between hyperglycemia and outcomes in 2,471 patients admitted to the hospital with community-acquired pneumonia. *Diabetes Care.* 2005;28:810-815.

Mehta SR, Yusuf S, Diaz R; CREATE-ECLA Trial Group Investigators. Effect of glucose-insulin-potassium infusion on mortality in patients with acute ST-segment elevation myocardial infarction: the CREATE-ECLA randomized controlled trial. *JAMA.* 2005;293:437-446.

Moghissi ES, Hirsch IB. Hospital management of diabetes. *Endocrinol Metab Clin North Am.* 2005;34:99-116.

Pinto DS, Skolnick AH, Kirtane AJ, et al. MU-shaped relationship of blood glucose with adverse outcomes among patients with ST-segment elevation myocardial infarction. *J Am Coll Cardiol.* 2005;46:178-80.

Rady MY, Johnson DJ, Patel BM, et al. Influence of individual characteristics on outcome of glycemic control in intensive care unit patients with or without diabetes mellitus. *Mayo Clin Proc.* 2005;80:1558-1567.

Smith WD, Winterstein AG, Johns T, et al. Causes of hyperglycemia and hypoglycemia in adult inpatients. *Am J Health Syst Pharm.* 2005;62:714-9.

Sung J, Bochicchio GV, Joshi M, et al. Admission hyperglycemia is predictive of outcome in critically ill trauma patients. *J Trauma.* 2005;59:80-83.

Svensson AM, McGuire DK, Abrahamsson P, et al. Association between hyper- and hypoglycaemia and 2 year all-cause mortality risk in diabetic patients with acute

coronary events. *Eur Heart J* 2005;26:1255-61.

Van den Berghe G, Schoonheydt K, Becx P, et al. Insulin therapy protects the central and peripheral nervous system of intensive care patients. *Neurology*. 2005;64:1348-1353.

2004

American College of Endocrinology Task Force on Inpatient Diabetes and Metabolic Control. ACE Position Statement on Inpatient Diabetes and Metabolic Control. *Endocr Pract*. 2004;10:77-82.

Antman EM, Anbe DT, Armstrong PW, et al. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction--executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 1999 Guidelines for the Management of Patients With Acute Myocardial Infarction). *Circulation*. 2004;110:588-636.

Bartnik M, Ryden L, Ferrari R, et al. The prevalence of abnormal glucose regulation in patients with coronary artery disease across Europe. The Euro Heart Survey on diabetes and the heart. *Eur Heart J*. 2004;25:1880-1890.

Braithwaite SS, Buie MM, Thompson CL, et al. Hospital hypoglycemia: not only treatment but also prevention. *Endocr Pract*. 2004;Mar-Apr;10 Suppl 2:89-99.

Clement S, Braithwaite SS, Magee MF, the American Diabetes Association Diabetes in Hospitals Writing Committee. Management of diabetes and hyperglycemia in hospitals. *Diabetes Care* 2004;27:553-591.

Cryer PE. Diverse causes of hypoglycemia-associated autonomic failure in diabetes. *N Engl J Med*. 2004;350:2272-9.

Davidson JA, Wilkinson A; International Expert Panel on New-Onset Diabetes after Transplantation. New-Onset Diabetes After Transplantation 2003 International Consensus Guidelines: an endocrinologist's view. *Diabetes Care*. 2004;27:805-12.

Furnary AP, Wu Y, Bookin SO. Effect of hyperglycemia and continuous intravenous insulin infusions on outcomes of cardiac surgical procedures: the Portland Diabetic Project. *Endocr Pract*. 2004;10:21-33.

Garber AJ, Seidel J, Armbruster M. Current standards of care for inpatient glycemic management and metabolic control: is it time for definite standards and targets? *Endocr Pract*. 2004;10 Suppl 2:10-2.

Goldberg PA, Siegel MD, Russell RR, et al. Experience with the continuous glucose monitoring system in a medical intensive care unit. *Diabetes Technol Ther*. 2004;6:339-47.

Goldberg PA, Siegel MD, Sherwin RS, et al. Implementation of a safe and effective insulin infusion protocol in a medical intensive care unit. *Diabetes Care*. 2004;27:461-7.

Grey NJ, Perdrizet GA. Reduction of nosocomial infections in the surgical intensive-care unit by

strict glycemic control. *Endocr Pract.* 2004;10 Suppl 2:46-52.

Hellman R. A systems approach to reducing errors in insulin therapy in the inpatient setting. *Endocr Pract.* 2004;10(Suppl 2):100-108.

Hirsch IB. Effect of insulin therapy on nonglycemic variables during acute illness. *Endocr Pract.* 2004;10 Suppl 2:63-70.

Krinsley JS. Effect of intensive glucose management protocol on the mortality of critically ill adult patients. *Mayo Clin Proc.* 2004;79:992-1000.

Laird AM, Miller PR, Kilgo PD, et al. Relationship of early hyperglycemia to mortality in trauma patients. *J Trauma.* 2004;56:1058-62.

Lazar HL, Chipkin SR, Fitzgerald CA, et al. Tight glycemic control in diabetic coronary artery bypass graft patients improves perioperative outcomes and decreases recurrent ischemic events. *Circulation.* 2004;109:1497-1502.

Levetan CS. Effect of hyperglycemia on stroke outcomes. *Endocr Pract.* 2004;10 Suppl 2:34-9.

Malmberg K. Role of insulin-glucose infusion in outcomes after acute myocardial infarction: the diabetes and insulin-glucose infusion in acute myocardial infarction (DIGAMI) study. *Endocr Pract.* 2004;10 Suppl 2:13-6.

Moghissi E. Hospital Management of Diabetes "Beyond the Sliding Scale." *Cleve Clin J Med.* 2004;71:801-8.

Norhammar A, Malmberg K, Diderholm E, et al. Diabetes mellitus: the major risk factor in unstable coronary artery disease even after consideration of the extent of coronary artery disease and benefits of revascularization. *J Am Coll Cardiol.* 2004;43:585-591.

Nylen ES, Muller B. Endocrine changes in critical illness. *J Intensive Care Med.* 2004;19:67-82.

Pittas AG, Siegel RD, Lau J. Insulin therapy for critically ill hospitalized patients: a meta-analysis of randomized, control trials. *Arch Inter Med.* 2004;164:2005-2011.

Timmer JR, van der Horst IC, Ottervanger JP, et al. Prognostic value of admission glucose in non-diabetic patients with myocardial infarction. *Am Heart J.* 2004;148:399-404.

Van den Berghe GH. Role of intravenous insulin therapy in critically ill patients. *Endocr Pract.* 2004;10 Suppl 2:17-20.

Vogelzang M, van der Horst ICC, Nijsten MW. Hyperglycemic index as a tool to assess glucose control: a retrospective study. *Crit Care;* 2004;8:R122-R127.

2003

Bates DW, Gawande AA. Improving patient safety with information technology. *N Engl J Med.* 2003;348:2526-2534.

Cryer PE, Davis SN, Shamooh H. Hypoglycemia in diabetes. *Diabetes Care*. 2003;26:1902-12.

Estrada CA, Young JA, Nifong LW, et al. Outcomes and perioperative hyperglycemia in patients with or without diabetes mellitus undergoing coronary artery bypass grafting. *Ann Thorac Surg*. 2003;75:1392-1399.

Finney SJ, Zekveld C, Elia A, et al. Glucose control and mortality in critically ill patients. *JAMA* 2003;290:2041-7.

Forster AJ, Murff HJ, Peterson JF, et al. The incidence and severity of adverse events affecting patients after discharge from the hospital. *Ann Intern Med*. 2003;138:161-7.

Furnary AP, Gao G, Grunkemeier GL, et al. Continuous insulin infusion reduces mortality in patients with diabetes undergoing coronary artery bypass grafting. *J Thorac Cardiovasc Surg*. 2003;125:1007-1021.

Krinsley JS. Association between hyperglycemia and increased hospital mortality in a heterogeneous population of critically ill patients. *Mayo Clin Proc*. 2003;78:1471-1478.

Lewis EF, Moye LA, Rouleau JL; CARE Study. Predictors of late development of heart failure in stable survivors of myocardial infarction: the CARE study. *J Am Coll Cardiol*. 2003;42:1446-1453.

Norhammar A, Malmberg K, Ryden L; Register of Information and Knowledge about Swedish Heart Intensive Care Admission (RIKS-HIA). Under utilisation of evidence-based treatment partially explains for the unfavourable prognosis in diabetic patients with acute myocardial infarction. *Eur Heart J*. 2003;24:838-844.

Trence DL, Kelly JL, Hirsch IB. The rationale and management of hyperglycemia for in-patients with cardiovascular disease: time for change. *J Clin Endocrinol Metab*. 2003;88:2430-2437.

Van den Berghe G, Wouters PJ, Bouillon R, et al. Outcome benefit of intensive insulin therapy in the critically ill: Insulin dose versus glycemic control. *Crit Care Med*. 2003;31:359-366.

Yendamuri S, Fulda GJ, Tinkoff GH. Admission hyperglycemia as a prognostic indicator in trauma. *J Trauma*. 2003;55:33-38.

2002

Bhattacharyya A, Christodoulides C, Kaushal K, et al. In-patient management of diabetes mellitus and patient satisfaction. *Diabet Med*. 2002;19:412-416.

Montori VM, Bistrian BR, McMahon MM. Hyperglycemia in acutely ill patients. *JAMA* 2002;288:2167-9.

Norhammar A, Tenerz A, Nilsson G, et al. Glucose metabolism in patients with acute myocardial infarction and no previous diagnosis of diabetes mellitus: a prospective study. *Lancet*. 2002;359:2140-2144.

Umpierrez GE, Isaacs SD, Bazargan N, et al. Hyperglycemia: an independent marker of in-

hospital mortality in patients with undiagnosed diabetes. *J Clin Endocrinol Metab.* 2002;87:978-982.

Wahab NN, Cowden EA, Pearce NJ; ICONS Investigators. Is blood glucose an independent predictor of mortality in acute myocardial infarction in the thrombolytic era? *J Am Coll Cardiol.* 2002;40:1748-1754.

2001

Bolk J, van der Ploeg T, Cornel JH, et al. Impaired glucose metabolism predicts mortality after a myocardial infarction. *Int J Cardiol.* 2001;79:207-214.

Boyd JC, Bruns DE. Quality specifications for glucose meters: assessment simulation modeling of errors in insulin dose. *Clin Chem.* 2001;47:209-14.

Capes SE, Hunt D, Malmberg K, et al. Stress hyperglycemia and prognosis of stroke in nondiabetic and diabetic patients: a systematic overview. *Stroke.* 2001;32:2426-2432.

Latham R, Lancaster AD, Covington JF, et al. The association of diabetes and glucose control with surgical site infection among cardiothoracic surgery patients. *Infect Control Hosp Epidemiol.* 2001;22:607-612.

Mukamal KJ, Nesto RW, Cohen MC, et al. Impact of diabetes on long-term survival after acute myocardial infarction: comparability of risk with prior myocardial infarction. *Diabetes Care.* 2001;24:1422-1427.

Pancorbo-Hidalgo PL, García-Fernandez FP, Ramírez-Pérez C. Complications associated with enteral nutrition by nasogastric tube in an internal medicine unit. *J Clin Nurs.* 2001;10:482-90.

Van Den Berghe G, Wouters P, Weekers F, et al. Intensive insulin therapy in critically ill patients. *N Engl J Med.* 2001;345:1359-1367.

2000

Capes SE, Hunt D, Malmberg K, et al. Stress hyperglycemia and increased risk of death after myocardial infarction in patients with and without diabetes: a systematic overview. *Lancet.* 2000;355:773-778.

1999

Furnary AP, Zerr KJ, Grunkemeier GL, et al. Continuous intravenous insulin infusion reduces the incidence of deep sternal wound infection in diabetic patients after cardiac surgical procedures. *Ann Thorac Surg.* 1999;67:352-360. Discussion 360-362.

Golden S, Peart-Vigilance C, Kao W, et al. Perioperative glycemic control and risk of infectious complications in a cohort of adults with diabetes. *Diabetes Care* 1999;22:1408-1414.

Malmberg K, Norhammar A, Wedel H, et al. Glycometabolic state at admission: important risk

marker of mortality in conventionally treated patients with diabetes mellitus and acute myocardial infarction: long-term results from the Diabetes and Insulin-Glucose Infusion in Acute Myocardial Infarction (DIGAMI) study. *Circulation*. 1999;99:2626-2632.

Norhammar AM, Ryden L, Malmberg K. Admission plasma glucose. Independent risk factor for long-term prognosis after myocardial infarction even in nondiabetic patients. *Diabetes Care* 1999;22:1827-31.

1998

Bates DW, Leape LL, Cullen DJ, et al. Effect of computerized physician order entry and a team intervention on prevention of serious medical errors. *JAMA*. 1998;280:1311-1316.

Cohen MR, Proulx SM, Crawford SY. Survey of hospital systems and common serious medication errors. *J Healthc Risk Manag*. 1998;18:16-27.

Levetan CS, Passaro M, Jablonski K, et al. Unrecognized diabetes among hospitalized patients. *Diabetes Care* 1998;21:246–249.

Pomposelli JJ, Baxter JK, 3rd, Babineau TJ et al. Early postoperative glucose control predicts nosocomial infection rate in diabetic patients. *JPEN J Parenter Enteral Nutr* 1998;22:77-81.

1997

Braunwald E. Shattuck lecture—cardiovascular medicine at the turn of the millennium: triumphs, concerns, and opportunities. *N Engl J Med*. 1997;337:1360-1369.

Malmberg K, DIGAMI Study Group. Prospective randomised study of intensive insulin treatment on long-term survival after acute myocardial infarction in patients with diabetes mellitus. *Br Med J* 1997;314:1512-5.

Queale WS, Seidler AJ, Brancati FL. Glycemic control and sliding scale insulin use in medical inpatients with diabetes mellitus. *Arch Intern Med*. 1997;157:545-52.

Zerr KJ, Furnary AP, Grunkemeier GL, et al. Glucose control lowers the risk of wound infection in diabetics after open heart operations. *Ann Thorac Surg*. 1997;63:356-361.

1995

Levetan CS, Salas JR, Wilets IF, et al. Impact of endocrine and diabetes team consultation on hospital length of stay for patients with diabetes. *Am J Med*. 1995;99:22-8.

Malmberg K, Ryden L, Efendic S, et al. Randomized trial of insulin-glucose infusion followed by subcutaneous insulin treatment in diabetic patients with acute myocardial infarction (DIGAMI study): effects on mortality at 1 year. *J Am Coll Cardiol*. 1995;26:57-65.

Mizock BA. Alterations in carbohydrate metabolism during stress: a review of the literature. *Am J Med* 1995;98:75-84.

1991

Mitrakou A, Ryan C, Veneman T, et al. Hierarchy of glycemic thresholds for counterregulatory hormone secretion, symptoms, and cerebral dysfunction. *Am J Physiol.* 1991;260(1 Pt 1):E67-74.

1988

Malmberg K, Ryden L. Myocardial infarction in patients with diabetes mellitus. *Eur Heart J.* 1988;9:259-264.