

# Clinical Evidence Supporting the Need for Glycemic Control in the Hospital Setting “The Case for Tight Glycemic Control in Hospitalized Patients”

## References and Key Readings

### Most Recent Consensus Statements

Moghissi ES, Korytkowski MT, DiNardo M; American Association of Clinical Endocrinologists; American Diabetes Association. American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycemic control. *Endocr Pract.* 2009;15:353-69.

Moghissi ES, Korytkowski MT, DiNardo M; American Association of Clinical Endocrinologists; American Diabetes Association. American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycemic control. *Diabetes Care.* 2009;32:1119-31.

Preiser JC, Devos P, Ruiz-Santana S, et al. A prospective randomised multi-centre controlled trial on tight glucose control by intensive insulin therapy in adult intensive care units: the Glucontrol study. *Intensive Care Med.* 2009;35:1738-48.

### 2008

Brunkenhorst FM, Engel C, Bloos F, for the German Competence Network Sepsis (SepNet). Intensive insulin therapy and pentastarch resuscitation in severe sepsis. *N Engl J Med.* 2008;358:125-39.

Cook CB, Jameson KA, Hartsell ZC, et al. Beliefs about hospital diabetes and perceived barriers to glucose management among inpatient midlevel practitioners. *Diabetes Educ.* 2008;34:75-83.

Kitabchi AE, Freire AX, Umpierrez GE. Evidence for strict inpatient blood glucose control: time to revise glycemic goals in hospitalized patients. *Metabolism.* 2008;57:116-20.

Kosiborod M, Inzucchi SE, Krumholz HM, et al. Glucometrics in patients hospitalized with acute myocardial infarction: defining the optimal outcomes-based measure of risk. *Circulation.* 2008;117:1018-27.

Matheny ME, Shubina M, Kimmel ZM, et al. Treatment intensification and blood glucose control among hospitalized diabetic patients. *J Gen Intern Med.* 2008;23:184-9.

Mills RD, Schwartz F, Shubrook JH. Evaluation of diabetes management in a rural community hospital. *Endocr Pract.* 2008;14:50-5.

Treggiari MM, Karir V, Yanez ND, et al. Intensive insulin therapy and mortality in critically ill patients. *Crit Care.* 2008;12(1):R29.

### 2007

Anthony M. Treatment of hypoglycemia in hospitalized adults: a descriptive study. *Diabetes Educ.* 2007;33:709-15.

Asudani D, Calles-Escandon J. Inpatient hyperglycemia: slide through the scale but cover the bases first. *J Hosp Med.* 2007;2 Suppl 1:23-32.

Becker T, Moldoveanu A, Cukierman T, et al. Clinical outcomes associated with the use of subcutaneous insulin-by-glucose sliding scales to manage hyperglycemia in hospitalized patients with pneumonia. *Diabetes Res Clin Pract.* 2007;78:392-7.

Cohen LS, Sedhom L, Salifu M, Friedman EA. Inpatient diabetes management: examining morning practice in an acute care setting. *Diabetes Educ.* 2007;33:483-92.

Cook CB, Castro JC, Schmidt RE, et al. Diabetes care in hospitalized noncritically ill patients: More evidence for clinical inertia and negative therapeutic momentum. *J Hosp Med.* 2007;2:203-11.

Cook CB, Moghissi E, Joshi R, et al. Inpatient point-of-care bedside glucose testing: preliminary data on use of connectivity informatics to measure hospital glycemic control. *Diabetes Technol Ther.* 2007;9:493-500.

Cook CB, McNaughton DA, Braddy CM, et al. Management of inpatient hyperglycemia: assessing perceptions and barriers to care among resident physicians. *Endocr Pract.* 2007;13:117-24.

Cook CB, Stockton L, Baird M, et al. Working to improve care of hospital hyperglycemia through statewide collaboration: the Georgia Hospital Association Diabetes Special Interest Group. *Endocr Pract.* 2007;13:45-50.

Diabetes Special Interest Group of the Partnership; for Health and Accountability. Tight blood glucose control in the hospital inpatient setting: a collaborative solution. *J Med Assoc Ga.* 2007;96:34-7, 39-40.

Elinav H, Wolf Z, Szalat A, et al. In-hospital treatment of hyperglycemia: effects of intensified subcutaneous insulin treatment. *Curr Med Res Opin.* 2007;23:757-65.

Gandhi GY, Nuttall GA, Abel MD, et al. Intensive intraoperative insulin therapy versus conventional glucose management during cardiac surgery: a randomized trial. *Ann Intern Med.* 2007;146:233-43.

Garg R, Bhutani H, Jarry A, et al. Provider response to insulin-induced hypoglycemia in hospitalized patients. *J Hosp Med.* 2007;2:258-60.

Gray CS, Hildreth AJ, Sandercock PA; GIST Trialists Collaboration. Glucose-potassium-insulin infusions in the management of post-stroke hyperglycaemia: the UK Glucose Insulin in Stroke Trial (GIST-UK). *Lancet Neurol.* 2007;6:397-406.

Hassan E. Hyperglycemia management in the hospital setting. *Am J Health Syst Pharm.* 2007;64(10 Suppl 6):S9-S14.

Held C, Gerstein HC, Yusuf S; ONTARGET/TRANSCEND Investigators. Glucose levels predict hospitalization for congestive heart failure in patients at high cardiovascular risk. *Circulation.* 2007;115:1371-5.

Hovorka R, Kremen J, Blaha J, et al. Blood glucose control by a model predictive control algorithm with variable sampling rate versus a routine glucose management protocol in cardiac surgery patients: a randomized controlled trial. *J Clin Endocrinol Metab.* 2007;92:2960-4.

Karon BS, Gandhi GY, Nuttall GA, et al. Accuracy of roche accu-chek inform whole blood capillary, arterial, and venous glucose values in patients receiving intensive intravenous insulin therapy after cardiac surgery. *Am J Clin Pathol.* 2007;127:919-26.

Kim S. Burden of hospitalizations primarily due to uncontrolled diabetes: implications of inadequate primary health care in the United States. *Diabetes Care.* 2007;30:1281-2.

Lansang MC, Harrell H. Knowledge on inpatient diabetes among fourth-year medical students. *Diabetes Care.* 2007;30:1088-91.

Rea RS, Donihi AC, Bobeck M, et al. Implementing an intravenous insulin infusion protocol in the intensive care unit. *Am J Health Syst Pharm.* 2007;64:385-95.

Reynolds LR, Cook AM, Lewis DA, et al. An institutional process to improve inpatient glycemic control. *Qual Manag Health Care.* 2007;16:239-49.

Umpierrez GE. Inpatient management of diabetes: an increasing challenge to the hospitalist physician. *J Hosp Med.* 2007;2 Suppl 1:33-5.

Umpierrez GE, Smiley D, Zisman A, et al. Randomized study of basal-bolus insulin therapy in the inpatient management of patients with type 2 diabetes (RABBIT 2 trial). *Diabetes Care.* 2007;30:2181-6.

Umpierrez GE, Palacio A, Smiley D. Sliding scale insulin use: myth or insanity? *Am J Med.* 2007;120:563-7.

Wexler DJ, Meigs JB, Cagliero E, et al. Prevalence of hyper- and hypoglycemia among inpatients with diabetes: a national survey of 44 U.S. hospitals. *Diabetes Care.* 2007;30:367-9.

Wheeler K, Crawford R, McAdams D, et al. Inpatient to outpatient transfer of diabetes care: perceptions of barriers to postdischarge followup in urban African American patients. *Ethn Dis.* 2007;17:238-43.

## **2006**

American College of Endocrinology Task Force on Inpatient Diabetes and Glycemic. ACE Position Statement on Inpatient Diabetes and Metabolic Control. *Endocr Pract.* 2006;12:3-13.

Baker EH, Janaway CH, Philips BJ, et al. Hyperglycaemia is associated with poor outcomes in patients admitted to hospital with acute exacerbations of chronic obstructive pulmonary disease. *Thorax.* 2006;61:284-289.

Cheung NW, Wong VW, McLean M. The Hyperglycemia: Intensive Insulin Infusion in Infarction (HI-5) study: a randomized controlled trial of insulin infusion therapy for myocardial infarction. *Diabetes Care.* 2006;29:765-770.

Donihi AC, Raval D, Saul M, et al. Prevalence and predictors of corticosteroid-related hyperglycemia in hospitalized patients. *Endocr Pract.* 2006;12:358-362.

Fonseca V. Newly diagnosed diabetes/hyperglycemia in hospitals: what should we do? *Endocr Pract.* 2006;12 Suppl 3:108-11.

Frontera JA, Fernandez A, Claassen J, et al. Hyperglycemia after SAH: predictors, associated complications, and impact on outcome. *Stroke.* 2006;37:199-203.

Furnary AP, Wu Y. Clinical effects of hyperglycemia in the cardiac surgery population: the Portland Diabetic Project. *Endocr Pract.* 2006;12 Suppl 3:22-6.

Green Conaway DL, Enriquez JR, Barberena JE, et al. Assessment of and physician response to glycemic control in diabetic patients presenting with an acute coronary syndrome. *Am Heart J.* 2006;152:1022-7.

Hirsch IB. Inpatient diabetes: review of data from the cardiac care unit. *Endocr Pract.* 2006;12 Suppl 3:27-34.

Knecht LA, Gauthier SM, Castro JC, et al. Diabetes care in the hospital: is there clinical inertia? *J Hosp Med.* 2006;1:151-60.

Korytkowski M, DiNardo M, Donihi AC, et al. Evolution of a diabetes inpatient safety committee. *Endocr Pract.* 2006;Suppl 3:91-99.

Olson L, Muchmore J, Lawrence CB. The benefits of inpatient diabetes care: improving quality of care and the bottom line. *Endocr Pract.* 2006;12 Suppl 3:35-42.

Schmeltz LR, Desantis AJ, Schmidt K, et al. Conversion of intravenous insulin infusions to subcutaneously administered insulin glargine in patients with hyperglycemia. *Endocr Pract.* 2006;12:641-650.

Umpierrez G, Maynard G. Glycemic chaos (not glycemic control) still the rule for inpatient care: how do we stop the insanity? *J Hosp Med.* 2006;1:141-4.

Van den Berghe G, Wilmer A, Hermans G, et al. Intensive insulin therapy in the medical ICU. *N Engl J Med.* 2006;354:449-461.

Van den Berghe G, Wouters PJ, Kesteloot K, et al. Analysis of healthcare resource utilization with intensive insulin therapy in critically ill patients. *Crit Care Med.* 2006;34:612-616.

Vanhorebeek I, Langouche L, Van den Berghe G. Intensive insulin therapy in the intensive care unit: update on clinical impact and mechanisms of action. *Endocr Pract.* 2006;12 Suppl 3:14-22.

## **2005**

Ainla T, Baburin A, Teesalu R, et al. The association between hyperglycaemia on admission and 180-day mortality in acute myocardial infarction patients with and without diabetes. *Diabet Med.* 2005;22:1321-1325.

Bochicchio GV, Sung J, Joshi M, et al. Persistent hyperglycemia is predictive of outcome in critically ill trauma patients. *J Trauma*. 2005;58:921-924.

Cheung NW, Napier B, Zaccaria C, et al. Hyperglycemia is associated with adverse outcomes in patients receiving total parenteral nutrition. *Diabetes Care*. 2005;28:2367-2371.

Conaway DG, O'Keefe JH, Reid KJ, et al. Frequency of undiagnosed diabetes mellitus in patients with acute coronary syndrome. *Am J Cardiol*. 2005;96:363-365.

Ishihara M, Kojima S, Sakamoto T; Japanese Acute Coronary Syndrome Study Investigators. Acute hyperglycemia is associated with adverse outcome after acute myocardial infarction in the coronary intervention era. *Am Heart J*. 2005;150:814-820.

Kosiborod M, Rathore SS, Inzucchi S et al. Admission glucose and mortality in elderly patients hospitalized with acute myocardial infarction: implications for patients with and without recognized diabetes. *Circulation*. 2005;111:3078-3086.

Lukins MB, Manninen PH. Hyperglycemia in patients administered dexamethasone for craniotomy. *Anesthesia & Analgesia*. 2005;100:1129-1133.

Malmberg K, Ryden L, Wedel H; DIGAMI 2 Investigators. Intense metabolic control by means of insulin in patients with diabetes mellitus and acute myocardial infarction (DIGAMI 2): effects on mortality and morbidity. *Eur Heart J*. 2005;26:650-661.

McAlister FA, Majumdar SR, Blitz S, et al. The relation between hyperglycemia and outcomes in 2,471 patients admitted to the hospital with community-acquired pneumonia. *Diabetes Care*. 2005;28:810-815.

Mehta SR, Yusuf S, Diaz R; CREATE-ECLA Trial Group Investigators. Effect of glucose-insulin-potassium infusion on mortality in patients with acute ST-segment elevation myocardial infarction: the CREATE-ECLA randomized controlled trial. *JAMA*. 2005;293:437-446.

Moghissi ES, Hirsch IB. Hospital management of diabetes. *Endocrinol Metab Clin North Am*. 2005;34:99-116.

Rady MY, Johnson DJ, Patel BM, et al. Influence of individual characteristics on outcome of glycemic control in intensive care unit patients with or without diabetes mellitus. *Mayo Clin Proc*. 2005;80:1558-1567.

Van den Berghe G, Schoonheydt K, Bex P, et al. Insulin therapy protects the central and peripheral nervous system of intensive care patients. *Neurology*. 2005;64:1348-1353.

## **2004**

American College of Endocrinology Task Force on Inpatient Diabetes and Metabolic Control. ACE Position Statement on Inpatient Diabetes and Metabolic Control. *Endocr Pract*. 2004;10:77-82.

Antman EM, Anbe DT, Armstrong PW, et al. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction--executive summary: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing

Committee to Revise the 1999 Guidelines for the Management of Patients With Acute Myocardial Infarction). *Circulation*. 2004;110:588-636.

Bartnik M, Ryden L, Ferrari R, et al. The prevalence of abnormal glucose regulation in patients with coronary artery disease across Europe. The Euro Heart Survey on diabetes and the heart. *Eur Heart J*. 2004;25:1880-1890.

Furnary AP, Wu Y, Bookin SO. Effect of hyperglycemia and continuous intravenous insulin infusions on outcomes of cardiac surgical procedures: the Portland Diabetic Project. *Endocr Pract*. 2004;10:21-33.

Garber AJ, Seidel J, Armbruster M. Current standards of care for inpatient glycemic management and metabolic control: is it time for definite standards and targets? *Endocr Pract*. 2004;10 Suppl 2:10-2.

Grey NJ, Perdrizet GA. Reduction of nosocomial infections in the surgical intensive-care unit by strict glycemic control. *Endocr Pract*. 2004;10 Suppl 2:46-52.

Hirsch IB. Effect of insulin therapy on nonglycemic variables during acute illness. *Endocr Pract*. 2004;10 Suppl 2:63-70.

Krinsley JS. Effect of intensive glucose management protocol on the mortality of critically ill adult patients. *Mayo Clin Proc*. 2004;79:992-1000.

Lazar HL, Chipkin SR, Fitzgerald CA, et al. Tight glycemic control in diabetic coronary artery bypass graft patients improves perioperative outcomes and decreases recurrent ischemic events. *Circulation*. 2004;109:1497-1502.

Levetan CS. Effect of hyperglycemia on stroke outcomes. *Endocr Pract*. 2004;10 Suppl 2:34-9.

Moghissi E. Hospital Management of Diabetes "Beyond the Sliding Scale." *Cleve Clin J Med*. 2004;71:801-8.

Norhammar A, Malmberg K, Diderholm E, et al. Diabetes mellitus: the major risk factor in unstable coronary artery disease even after consideration of the extent of coronary artery disease and benefits of revascularization. *J Am Coll Cardiol*. 2004;43:585-591.

Pittas AG, Siegel RD, Lau J. Insulin therapy for critically ill hospitalized patients: a Meta analysis of randomized, control trials. *Arch Intern Med*. 2004;164:2005-2011.

Timmer JR, van der Horst IC, Ottervanger JP, et al. Prognostic value of admission glucose in non-diabetic patients with myocardial infarction. *Am Heart J*. 2004;148:399-404.

Van den Berghe GH. Role of intravenous insulin therapy in critically ill patients. *Endocr Pract*. 2004;10 Suppl 2:17-20.

### **2003**

Estrada CA, Young JA, Nifong LW, et al. Outcomes and perioperative hyperglycemia in patients with or without diabetes mellitus undergoing coronary artery bypass grafting. *Ann Thorac Surg*. 2003;75:1392-1399.

Furnary AP, Gao G, Grunkemeier GL, et al. Continuous insulin infusion reduces mortality in patients with diabetes undergoing coronary artery bypass grafting. *J Thorac Cardiovasc Surg.* 2003;125:1007-1021.

Krinsley JS. Association between hyperglycemia and increased hospital mortality in a heterogeneous population of critically ill patients. *Mayo Clin Proc.* 2003;78:1471-1478.

Lewis EF, Moye LA, Rouleau JL; CARE Study. Predictors of late development of heart failure in stable survivors of myocardial infarction: the CARE study. *J Am Coll Cardiol.* 2003;42:1446-1453.

Norhammar A, Malmberg K, Ryden L; Register of Information and Knowledge about Swedish Heart Intensive Care Admission (RIKS-HIA). Under utilisation of evidence based treatment partially explains for the unfavourable prognosis in diabetic patients with acute myocardial infarction. *Eur Heart J.* 2003;24:838-844.

Trence DL, Kelly JL, Hirsch IB. The rationale and management of hyperglycemia for in-patients with cardiovascular disease: time for change. *J Clin Endocrinol Metab.* 2003;88:2430-2437.

Van den Berghe G, Wouters PJ, Bouillon R, et al. Outcome benefit of intensive insulin therapy in the critically ill: Insulin dose versus glycemic control. *Crit Care Med.* 2003;31:359-366.

Yendamuri S, Fulda GJ, Tinkoff GH. Admission hyperglycemia as a prognostic indicator in trauma. *J Trauma.* 2003;55:33-38.

## **2002**

Bhattacharyya A, Christodoulides C, Kaushal K, et al. In-patient management of diabetes mellitus and patient satisfaction. *Diabet Med.* 2002;19:412-416.

Norhammar A, Tenerz A, Nilsson G, et al. Glucose metabolism in patients with acute myocardial infarction and no previous diagnosis of diabetes mellitus: a prospective study. *Lancet.* 2002;359:2140-2144.

Sala J, Masia R, Gonzalez de Molina F-J, et al. Short-term mortality of myocardial infarction patients with diabetes or hyperglycemia during admission. *J Epidemiol Community Health.* 2002;56:707-712.

Umpierrez GE, Isaacs SD, Bazargan N, et al. Hyperglycemia: an independent marker of in-hospital mortality in patients with undiagnosed diabetes. *J Clin Endocrinol Metab.* 2002;87:978-982.

Wahab NN, Cowden EA, Pearce NJ; ICONS Investigators. Is blood glucose an independent predictor of mortality in acute myocardial infarction in the thrombolytic era? *J Am Coll Cardiol.* 2002;40:1748-1754.

## **2001**

Bolk J, van der Ploeg T, Cornel JH, et al. Impaired glucose metabolism predicts mortality after a myocardial infarction. *Int J Cardiol.* 2001;79:207-214.

Capes SE, Hunt D, Malmberg K, et al. Stress hyperglycemia and prognosis of stroke in non diabetic and diabetic patients: a systematic overview. *Stroke*. 2001;32:2426-2432.

Latham R, Lancaster AD, Covington JF, et al. The association of diabetes and glucose control with surgical site infection among cardiothoracic surgery patients. *Infect Control Hosp Epidemiol*. 2001;22:607-612.

Mukamal KJ, Nesto RW, Cohen MC, et al. Impact of diabetes on long-term survival after acute myocardial infarction: comparability of risk with prior myocardial infarction. *Diabetes Care*. 2001;24:1422-1427.

Van Den Berghe G, Wouters P, Weekers F, et al. Intensive insulin therapy in critically ill patients. *N Engl J Med*. 2001;345:1359-1367.

## **2000**

Capes SE, Hunt D, Malmberg K, et al. Stress hyperglycemia and increased risk of death after myocardial infarction in patients with and without diabetes: a systematic overview. *Lancet*. 2000;355:773-778.

## **1999**

Furnary AP, Zerr KJ, Grunkemeier GL, et al. Continuous intravenous insulin infusion reduces the incidence of deep sternal wound infection in diabetic patients after cardiac surgical procedures. *Ann Thorac Surg*. 1999;67:352-360. Discussion 360-362.

Malmberg K, Norhammar A, Wedel H, et al. Glycometabolic state at admission: important risk marker of mortality in conventionally treated patients with diabetes mellitus and acute myocardial infarction: long-term results from the Diabetes and Insulin-Glucose Infusion in Acute Myocardial Infarction (DIGAMI) study. *Circulation*. 1999;99:2626-2632.

## **1998**

Pomposelli JJ, Baxter JK, 3rd, Babineau TJ et al. Early postoperative glucose control predicts nosocomial infection rate in diabetic patients. *JPEN J Parenter Enteral Nutr* 1998;22:77-81.

## **1997**

Braunwald E. Shattuck lecture—cardiovascular medicine at the turn of the millennium: triumphs, concerns, and opportunities. *N Engl J Med*. 1997;337:1360-1369.

Malmberg K, DIGAMI Study Group. Prospective randomised study of intensive insulin treatment on long-term survival after acute myocardial infarction in patients with diabetes mellitus. *Br Med J* 1997;314:1512-5.

Zerr KJ, Furnary AP, Grunkemeier GL, et al. Glucose control lowers the risk of wound infection in diabetics after open heart operations. *Ann Thorac Surg*. 1997;63:356-361.

## **1995**

Malmberg K, Ryden L, Efendic S, et al. Randomized trial of insulin-glucose infusion followed by subcutaneous insulin treatment in diabetic patients with acute myocardial infarction (DIGAMI study): effects on mortality at 1 year. *J Am Coll Cardiol.* 1995;26:57-65.

**1988**

Malmberg K, Ryden L. Myocardial infarction in patients with diabetes mellitus. *Eur Heart J.* 1988;9:259-264.