

Successful Models of Implementation

References and Key Readings

Most Recent Consensus Statements

Moghissi ES, Korytkowski MT, DiNardo M; American Association of Clinical Endocrinologists; American Diabetes Association. American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycemic control. *Endocr Pract.* 2009;15:353-69.

Moghissi ES, Korytkowski MT, DiNardo M; American Association of Clinical Endocrinologists; American Diabetes Association. American Association of Clinical Endocrinologists and American Diabetes Association consensus statement on inpatient glycemic control. *Diabetes Care.* 2009;32:1119-31.

2009

Falciglia M, Freyberg RW, Almenoff PL, et al. Hyperglycemia-related mortality in critically ill patients varies with admission diagnosis. *Crit Care Med.* 2009;37:3001-9.

Umpierrez GE. Basal versus sliding-scale regular insulin in hospitalized patients with hyperglycemia during enteral nutrition therapy. *Diabetes Care* 2009; 32:594–596.

Umpierrez GE, Hor T, Smiley D, et al. Comparison of Inpatient Insulin Regimens with Detemir plus Aspart Versus Neutral Protamine Hagedorn plus Regular in Medical Patients with Type 2 Diabetes. *J Clin Endocrinol Metab.* 2009;94:564-9.

2008

Matheny ME, Shubina M, Kimmel ZM, et al. Treatment intensification and blood glucose control among hospitalized diabetic patients. *J Gen Intern Med.* 2008;23:184-9.

Mills RD, Schwartz F, Shubrook JH. Evaluation of diabetes management in a rural community hospital. *Endocr Pract.* 2008;14:50-5.

Trujillo JM, Barsky EE, Greenwood BC, et al. Improving glycemic control in medical inpatients: a pilot study. *J Hosp Med.* 2008;3:55-63.

2007

Asudani D, Calles-Escandon J. Inpatient hyperglycemia: slide through the scale but cover the bases first. *J Hosp Med.* 2007;2 Suppl 1:23-32.

Becker T, Moldoveanu A, Cukierman T, et al. Clinical outcomes associated with the use of subcutaneous insulin-by-glucose sliding scales to manage hyperglycemia in hospitalized patients with pneumonia. *Diabetes Res Clin Pract.* 2007;78:392-7.

Cook CB, Castro JC, Schmidt RE, Gauthier et al. Diabetes care in hospitalized noncritically ill patients: More evidence for clinical inertia and negative therapeutic momentum. *J Hosp Med.* 2007;2:203-11.

Cook CB, McNaughton DA, Braddy CM, et al. Management of inpatient hyperglycemia: Asssing perceptions and barriers to care among resident physicians. *Endocr Pract* 2007;13:117-125.

Cook CB, Stockton L, Baird M, et al. Working to improve care of hospital hyperglycemia through statewide collaboration: the Georgia Hospital Association Diabetes Special Interest Group. *Endocr Pract.* 2007;13:45-50.

Krinsley JS. Translating evidence into practice in managing inpatient hyperglycemia. *J Hosp Med.* 2007;2:13-19.

Lansang MC, Harrell H. Knowledge on inpatient diabetes among fourth-year medical students. *Diabetes Care.* 2007;30:1088-91.

Reynolds LR, Cook AM, Lewis DA, et al. An institutional process to improve inpatient glycemic control. *Qual Manag Health Care.* 2007;16:239-49.

Umpierrez GE, Smiley D, Zisman A, et al. Randomized study of basal-bolus insulin therapy in the inpatient management of patients with type 2 diabetes (RABBIT 2 trial). *Diabetes Care.* 2007;30:2181-6.

Umpierrez GE, Palacio A, Smiley D. Sliding scale insulin use: myth or insanity? *Am J Med.* 2007;120:563-7.

2006

ACE/ADA Task Force on Inpatient Diabetes. American College of Endocrinology and American Diabetes Association consensus statement on inpatient diabetes and glycemic control. *Endocr Pract.* 2006;12:458-468.

Alfonso A, Koops MK, Mong DP, Vigersky RA. Glycemic control with regular versus lispro insulin sliding scales in hospitalized Type 2 diabetics. *J Diabetes Complications.* 2006 May-Jun;20(3):153-7.

DeSantis AJ, Schmeltz LR, Schmidt K, et al. Inpatient management of hyperglycemia: the Northwestern experience. *Endocr Pract.* 2006;12:491-505.

Donaldson S, Villanueva G, Rondinelli L, et al. Rush University guidelines and protocols for the management of hyperglycemia in hospitalized patients: elimination of the sliding scale and improvement of glycemic control throughout the hospital. *Diabetes Educ.* 2006;32:954-962.

Donihi AC, DiNardo MM, DeVita MA, Korytkowski MT. Use of a standardized protocol to decrease medication errors and adverse events related to sliding scale insulin. *Qual Saf Health Care.* 2006;15:89-91.

Fraser DD, Robley LR, Ballard NM, Peno-Green LA. Collaborative development of an insulin nomogram for intensive insulin therapy. *Crit Care Nurs Q.* 2006;29:96-105.

Golightly LK, Jones MA, Hamamura DH, et al. Management of diabetes mellitus in hospitalized patients: efficiency and effectiveness of sliding-scale insulin therapy. *Pharmacotherapy.* 2006 Oct;26(10):1421-32.

Mozdzan M, Ruxer J, Loba J, Siejka A, Markuszewski L. Safety of various methods of intensive insulin therapy in hospital condition assessed by hypoglycaemic episodes detected with the use of continuous glucose monitoring system. *Adv Med Sci.* 2006;51:133-6.

Newton CA, Young S. Financial implications of glycemic control: results of an inpatient diabetes management program. *Endocr Pract.* 2006;12(suppl 3):43-48.

Olson L, Muchmore J, Lawrence CB. The benefits of inpatient diabetes care: improving quality of care and the bottom line. *Endocr Pract.* 2006;12(suppl 3):35-42.

Osburne RC, Cook CB, Stockton L, et al. Improving hyperglycemia management in the intensive care unit: preliminary report of a nurse-driven quality improvement project using a redesigned insulin infusion algorithm. *Diabetes Educ.* 2006;32:394-403.

Schmeltz LR, Desantis AJ, Schmidt K, et al. Conversion of intravenous insulin infusions to subcutaneously administered insulin glargine in patients with hyperglycemia. *Endocr Pract.* 2006;12:641-650.

Schnipper JL, Barsky EE, Shaykevich S, et al. Inpatient management of diabetes and hyperglycemia among general medicine patients at a large teaching hospital. *J Hosp Med.* 2006;1:145-50.

2005

Ku SY, Sayre CA, Hirsch IB, et al. New insulin infusion protocol Improves blood glucose control in hospitalized patients without increasing hypoglycemia. *Jt Comm J Qual Patient Saf.* 2005;31:141-147.

Moghissi ES, Hirsch IB. Hospital management of diabetes. *Endocrinol Metab Clin North Am.* 2005;34:99-116.

2004

American College of Endocrinology Task Force on Inpatient Diabetes and Metabolic Control. ACE Position Statement on Inpatient Diabetes and Metabolic Control. *Endocr Pract.* 2004;10:77-82.

Goldberg PA, Siegel MD, Russell RR, et al. Experience with the continuous glucose monitoring system in a medical intensive care unit. *Diabetes Technol Ther.* 2004;6:339-347.

Hellman R. A systems approach to reducing errors in insulin therapy in the inpatient setting. *Endocr Pract.* 2004;10(suppl 2):100-108.

Magee MF, Clement S. Subcutaneous insulin therapy in the hospital setting: issues, concerns, and implementation. *Endocr Pract.* 2004;10(suppl 2):81-88.

2003

Krinsley JS. Association between hyperglycemia and increased hospital mortality in a heterogeneous population of critically ill patients. *Mayo Clin Proc.* 2003;78:1471-1478.

2002

Markovitz LJ, Wiechmann RJ, Harris N, et al. Description and evaluation of a glycemic management protocol for patients with diabetes undergoing heart surgery. *Endocr Pract.* 2002;8:10-18.

1998

Levetan CS, Passaro M, Jablonski K, et al. Unrecognized diabetes among hospitalized patients. *Diabetes Care.* 1998;21:246-249.

1997

Queale WS, Seidler AJ, Brancati FL. Glycemic control and sliding scale insulin use in medical inpatients with diabetes mellitus. *Arch Intern Med.* 1997;157:545-52.